

Native Burkholderia sp. Lipoprotein Lipase

Product Information

Cat#	NATE-0417
Similar	LPL
Source	Burkholderia sp.
Description	Lipoprotein lipase (LPL) (EC 3.1.1.34) is a member of the lipase gene family, which includes pancreatic lipase, hepatic lipase, and endothelial lipase. It is a water-soluble enzyme that hydrolyzes triglycerides in lipoproteins, such as those found in chylomicrons and very low-density lipoproteins (VLDL), into two free fatty acids and one monoacylglycerol molecule. It is also involved in promoting the cellular uptake of chylomicron remnants, cholesterol-rich lipoproteins, and free fatty acids. LPL requires ApoC-II as a cofactor. LPL is attached to the luminal surface of endothelial cells in capillaries by the protein glycosylphosphatidylinositol HDL-binding protein 1 (GPIHBP1) and by heparin sulfated proteoglycans. It is most widely distributed in adipose, heart, and skeletal muscle tissue, as well as in lactating mammary glands.
Form	lyophilized powder
Activity	> 50,000 units/mg solid
CAS No.	9004-02-8
Unit Definition	One unit will release 1.0 nmole of p-nitrophenol per min at pH 7.2 at 37°C using p-nitrophenyl butyrate as substrate.
Storage	?20°C
Synonyms	lipoprotein lipase; clearing factor lipase; diglyceride lipase; diacylglycerol lipase; postheparin esterase; diglyceride lipase; postheparin lipase; diacylglycerol hydrolase; lipemia-clearing factor; EC 3.1.1.34; 9004-02-8; LPL
Enzyme Commission Number	EC 3.1.1.34
Abbr	LPL, Native (Burkholderia sp.)



Creative Enzymes

Diagnostic Enzymes

Native Burkholderia sp. Lipoprotein Lipase

Alias	LPL
Applications	Lipoprotein lipase has been used in a study to assess the role of lipogenic enzymes in colorectal cancer. It has also been used in a study to investigate lipasemic activity of low molecular weight heparin in rats.
Product Overview	Lipoprotein lipase hydrolyzes triglycerides in plasma lipoproteins causing release of fatty acids for metabolic purposes in muscles and adipose tissue.