

DNase I (Glycerol-free)

Product Information

Cat#	DIA-469
Description	A deoxyribonuclease that randomly cleaves ssDNA and dsDNA, producing 5'-phosphate oligonucleotides. Activity depends on Ca ²⁺ and is activated by Mg ²⁺ or Mn ²⁺ . Glycerol-free, suitable for lyophilization.
Unit Definition	One unit (U) is defined as the amount of enzyme required to completely degrade 1 μ g of pBR322 plasmid DNA in 10 minutes at 37 °C.
Storage	–20 °C
Synonyms	Deoxyribonuclease I
Contaminants	Free of RNase contamination
Applications	Preparation of DNA-free RNA samples; removal of genomic DNA contamination prior to RT-PCR; elimination of DNA templates after in vitro transcription with T7, T3, or SP6 RNA polymerases; footprinting assays for DNA-protein interaction analysis; nick translation when combined with DNA polymerase I; DNA fragmentation to generate random libraries in the presence of Mn ²⁺ ; positive control for DNA fragmentation in apoptosis TUNEL assays.
Product Overview	Product Components: 200 U: DNase I (Glycerol-free, 1 U/μL): 200 μL 10x Reaction Buffer with MgSO ₄ : 1 mL

25 mM EDTA: 1 mL

2000 U:

DNase I (Glycerol-free, 1 U/µL): 2 mL 10× Reaction Buffer with MgSO₄: 10 mL

25 mM EDTA: 10 mL

20 KU:

DNase I (Glycerol-free, 1U/μl): 20 mL 10× Reaction Buffer with MgSO₄: 100 ml

Fax:1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA



DNase I (Glycerol-free)

25mM EDTA: 100 mL

50 KU:

DNase I (Glycerol-free, 1 U/μL): 50 mL

Package 200 U/2000 U/20 KU/50 KU

Fax:1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA