



## Native Bovine intestinal mucosa Alkaline Phosphatase

### Product Information

<b>Cat#</b>	NATE-0053
<b>Abbr</b>	ALP, Native (Bovine)
<b>Alias</b>	ALP; ALKP
<b>Similar</b>	ALP
<b>Species</b>	Bovine
<b>Source</b>	Bovine intestinal mucosa
<b>Description</b>	<p>Alkaline phosphatase (ALP, ALKP, ALPase, Alk Phos) (EC 3.1.3.1) is a hydrolase enzyme responsible for removing phosphate groups from many types of molecules, including nucleotides, proteins, and alkaloids. The process of removing the phosphate group is called dephosphorylation. As the name suggests, alkaline phosphatases are most effective in an alkaline environment. It is sometimes used synonymously as basic phosphatase.</p>
<b>Applications</b>	<p>Alkaline phosphatase can be used to dephosphorylate casein and other proteins. Alkaline phosphatase may be also be used to dephosphorylate the 5'-termini of DNA or RNA to prevent self-ligation. DNA or RNA can also be tagged with radiolabeled phosphate (via T4 polynucleotide kinase) after dephosphorylation with alkaline phosphatase. Alkaline phosphatase is used for conjugation to antibodies and other proteins for ELISA, Western blotting, and histochemical detection. It is routinely used to dephosphorylate proteins and nucleic acids. It may be used for protein labeling when high sensitivity is required. Alkaline phosphatase may be also be used to dephosphorylate the 5'-termini of DNA or RNA to prevent self-ligation. DNA or RNA can also be tagged with radiolabeled phosphate (via T4 polynucleotide kinase) after dephosphorylation with alkaline phosphatase. This product has been used to study the mon oclonal alkaline phosphatase-anti-alkaline phosphatase (APAAP) complex. High specific activity grade recommended for antibody and protein conjugation.</p>
<b>Product Overview</b>	<p>Bovine intestinal alkaline phosphatase is a dimeric, membrane-derived glycoprotein. At least three isoforms exist, which typically possess two N-linked and one or more O-</p>

## Native Bovine intestinal mucosa Alkaline Phosphatase

linked glycans per monomer.<sup>2</sup> The enzyme requires zinc, and magnesium or calcium divalent ions for activity. Bovine intestinal alkaline phosphatase is a dimeric, membrane-derived glycoprotein. At least three isoforms exist, which typically possess two N-linked and one or more O-linked glycans per monomer.

<b>Form</b>	Freeze dried powder
<b>Enzyme Commission Number</b>	EC 3.1.3.1
<b>Activity</b>	> 2,000 units/mg protein
<b>CAS No.</b>	9001-78-9
<b>Unit Definition</b>	One DEA unit will hydrolyze 1 $\mu$ mole of 4-nitrophenyl phosphate per minute at pH 9.8 at 37°C. (One glycine unit is equivalent to ~3 DEA units)
<b>Storage</b>	2-8°C
<b>Warnings</b>	Package sizes are based on DEA units
<b>Pathway</b>	Endochondral Ossification, organism-specific biosystem (from WikiPathways) Folate biosynthesis, organism-specific biosystem (from KEGG) Folate biosynthesis, conserved biosystem (from KEGG) Metabolic pathways, organism-specific biosystem (from KEGG) TNF-alpha NF-kB Signaling Pathway, organism-specific biosystem (from WikiPathways)
<b>Function</b>	The peri-partum characteristics of serum bone-specific alkaline phosphatase (BAP) and urinary deoxypyridinoline (DPD) in dairy cattle are reported. Results indicate that the presence of glycosylphosphatidylinositol increases the stability of alkaline phosphatase against chemical denaturation while it decreases its refolding yield by the artificial chaperone refolding technique. Reliable and reproducible estimates of $k$ (cat) and $K$ (m) from only two or three progress curves were obtained using alkaline phosphatase. GPI-anchored proteins are localized on the outer layer of plasma membranes and clustered in microdomains generally called lipid rafts.
<b>Synonyms</b>	Alkaline phosphatase; ALP; ALKP; ALPase; Alk Phos; EC 3.1.3.1; Alkaline



**Creative Enzymes**

*Diagnostic Enzymes*

## **Native Bovine intestinal mucosa Alkaline Phosphatase**

phosphomonoesterase; Glycerophosphatase; Phosphomonoesterase

---

Tel: 1-631-562-8517 1-516-512-3133

Email: [info@creative-enzymes.com](mailto:info@creative-enzymes.com)

Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY11967, USA