

## Native Sweet almond $\beta$ -Glucosidase

## **Product Information**

Cat#	DIA-195
Abbr	β-Glucosidase (Sweet almond)
Alias	elaterase
Similar	β-Glucosidase
Source	Sweet almond
Description	Beta-glucosidase is a glucosidase enzyme that acts upon β1->4 bonds linking two glucose or glucose-substituted molecules (i.e., the disaccharide cellobiose). It is one of the cellulases, enzymes involved in the decomposition of cellulose and related polysaccharides; more specifically, an exocellulase with specificity for a variety of beta-D-glycoside substrates. It catalyzes the hydrolysis of terminal non-reducing residues in beta-D-glucosides with release of glucose.
Applications	This enzyme is useful for structural investigations of carbohydrates and for the enzymatic determination of $\alpha$ -amylase when coupled with $\alpha$ -glucosidase in clinical analysis.
Appearance	Light yellow amorphous powder, lyophilized
Form	Freeze dried powder
Enzyme Commission Number	EC 3.2.1.21
Activity	10U/mg-solid or more (containing approx. 50% of BSA)
CAS No.	9001-22-3
Contaminants	α-Amylase < 5.0×10 <sup>-4</sup> %
Molecular Weight	approx. 110 kDa
Isoelectric point	7.3
pH Stability	pH 6.0-9.0 (25°C, 64hr)

Fax:1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA



## Native Sweet almond β-Glucosidase

2.8×10 <sup>-3</sup> M (p-Nitrophenyl-ß-D-glucopyranoside), 3.3×10 <sup>-3</sup> M (2,4-Dichlorophenyl-ß-D-glucopyranoside)
2 subunits per mol of enzyme
5.5
50-55°C
below 50°C (pH 7.3, 1hr)
Stable at-20°C for at least 6 months (A decrease in activity of ca. 10% may occur at 5°C within 6 months)
Bovine serum albumin (BSA), glutathione (reduced)
EC 3.2.1.21; gentiobiase; cellobiase; emulsin; elaterase; aryl-beta-glucosidase; beta-D-glucosidase; beta-glucoside glucohydrolase; arbutinase; amygdalinase; p-nitrophenyl beta-glucosidase; primeverosidase; amygdalase; linamarase; salicilinase; beta-1,6-glucosidase.

Fax:1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA