

Trypsin from Porcine, Recombinant

Product Information

Cat#	NATE-1148
Abbr	Trypsin, Recombinant (Porcine)
Similar	Trypsin
Source	Porcine
Description	<p>Trypsin (EC 3.4.21.4) is a serine protease from the PA clan superfamily, found in the digestive system of many vertebrates, where it hydrolyses proteins. Trypsin is produced in the pancreas as the inactive protease trypsinogen. Trypsin cleaves peptide chains mainly at the carboxyl side of the amino acids lysine or arginine, except when either is followed by proline. It is used for numerous biotechnological processes. The process is commonly referred to as trypsin proteolysis or trypsinisation, and proteins that have been digested/treated with trypsin are said to have been trypsinized.</p>
Applications	<p>Trypsin can be used to re-suspend cells adherent to the cell culture dish wall during the process of harvesting cells. Trypsin can also be used to dissociate dissected cells (for example, prior to cell fixing and sorting). Trypsin is commonly used in biological research during proteomics experiments to digest proteins into peptides for mass spectrometry analysis, e.g. in-gel digestion. Trypsin is particularly suited for this, since it has a very well defined specificity, as it hydrolyzes only the peptide bonds in which the carbonyl group is contributed either by an Arg or Lys residue. Trypsin can also be used to dissolve bloodclots in its microbial form and treat inflammation in its pancreatic form. During the industrial production of insulin, trypsin is necessary.</p>
Appearance	Colorless aqueous solution
Form	Freeze dried powder
Enzyme Commission Number	EC 3.4.21.4
Activity	120 Units/mg protein



Creative Enzymes

Diagnostic Enzymes

Trypsin from Porcine, Recombinant

Molecular Weight	24KDa (Determined by SDS-PAGE)
Purity	>90% (by SDS-PAGE)
Unit Definition	One unit will produce an increase by 0.18 per min at 247nm, at pH8.1, at 25°C, using TAME as substrate, Reaction volume = 3.0 mL (1 cm light path).
Storage	4°C, store at -20°C/-80°C for long-term preservation , Avoid multiple freeze-thaw cycles.
Buffer	20mM NaAc , pH3.5
Synonyms	α -trypsin; β -trypsin; cocoonase; parenzyme; parenzymol; tryptar; trypure; pseudotrypsin; tryptase; tripcellim; sperm receptor hydrolase; Alpha-trypsin; Beta-trypsin; EC 3.4.21.4; Trypsin; Acetyltrypsin