

Glutamate dehydrogenase, Recombinant

Product Information

| Cat# | NATE-1145 |
|------------------|--|
| Abbr | GLDH, Recombinant |
| Alias | GLDH |
| Similar | GLDH |
| Description | Glutamate dehydrogenase (GLDH) is an enzyme, present in most microbes and the mitochondria of eukaryotes, as are some of the other enzymes required for urea synthesis, that converts glutamate to α -ketoglutarate, and vice versa. In animals, the produced ammonia is usually used as a substrate in the urea cycle. Typically, the α -ketoglutarate to glutamate reaction does not occur in mammals, as glutamate dehydrogenase equilibrium favours the production of ammonia and α -ketoglutarate. |
| Applications | Except glutamate dehydrogenation, GLDH can also catalytic the deaminase of other amino acids such as L-valine, L-2-aminobutyric acid and L-leucine. The main measuring method is continuous monitoring. Moreover, GLDH catalyzes the reaction of α -ketoglutarate, H+,ammonia and NADH to generating glutamic. Since NADH is the color source of many biochemical assays, therefore the reaction catalyzed by the corresponding GLDH is widely used to detect the final step of biochemical detection reagent. |
| Appearance | White powder, lyophilized |
| Product Overview | Glutamate dehydrogenase (GLDH, EC 1.4.1.2) is the enzyme present in the mitochondrial matrix of the cell. It can convert glutamic acid to α-ketoglutarate and |

Glutamate dehydrogenase (GLDH, EC 1.4.1.2) is the enzyme present in the mitochondrial matrix of the cell. It can convert glutamic acid to α -ketoglutarate and catalyze the reverse reaction as well. GLDH is one of the allosteric enzymes which generated in the body through oxidative dehydrogenation, aminotransferase, combined dehydrogenation, non-oxidative dehydrogenation reaction and so on. The combined dehydrogenation is the most important reaction in the body. GLDH is an important molecular in the assimilation and alienation pathway. It is rich in the liver lobule cells followed by the kidney, pancreas, brain, small intestine and heart. Since GLDH is a kind of liver-specific enzyme, the change of GLDH activity in the peripheral blood reflects the

Fax:1-631-938-8127 45-1 Ramsey Road, Shirley, NY11967, USA



Glutamate dehydrogenase, Recombinant

| | changes of the liver function to some extent. |
|------------------|---|
| Form | Freeze dried powder |
| Enzyme | EC 1.4.1.2 |
| Commission | |
| Number | |
| Activity | >400U/mg |
| Molecular Weight | About 65kDa (SDS-PAGE detection) |
| Purity | >90% (SDS-PAGE test) |
| Unit Definition | One unit will convert 1µmol NADH per min at pH 8.3and at 37°C. |
| Storage | 4°C, store at -20°C for long-term preservation. |
| Buffer | 20mM Tris, PH8.0 |
| Synonyms | glutamate dehydrogenase; glutamic dehydrogenase; glutamate dehydrogenase |
| | (NAD+); glutamate oxidoreductase; glutamic acid dehydrogenase; L-glutamate |
| | dehydrogenase; NAD+-dependent glutamate dehydrogenase; NAD+-dependent |
| | glutamic dehydrogenase; NAD+-glutamate dehydrogenase; NAD+-linked glutamate |
| | dehydrogenase; NAD+-linked glutamic dehydrogenase; NAD+-specific glutamic |
| | dehydrogenase; NAD+-specific glutamate dehydrogenase; NAD+:glutamate |
| | oxidoreductase; NADH-linked glutamate dehydrogenase; GLDH; EC 1.4.1.2 |

Fax:1-631-938-8127 45-1 Ramsey Road, Shirley, NY11967, USA