

Native Microorganism Glucose Dehyrogenase (NAD(P)-dependent)

Product Information

Cat#	DIA-191
Abbr	Glucose Dehyrogenase (Microorganism)
Similar	Glucose Dehyrogenase
Source	Microorganism
Description	In enzymology, a glucose 1-dehydrogenase (EC 1.1.1.47) is an enzyme that catalyzes the chemical reaction: beta-D-glucose + NAD(P)+ ↔ D-glucono-1,5-lactone + NAD(P)H + H+. The 3 substrates of this enzyme are beta-D-glucose, NAD+, and NADP+, whereas its 4 products are D-glucono-1,5-lactone, NADH, NADPH, and H+. This enzyme belongs to the family of oxidoreductases, specifically those acting on the CH-OH group of donor with NAD+ or NADP+ as acceptor.
Applications	This enzyme is useful for enzymatic determination of D-Glucose.
Appearance	White amorphous powder, lyophilized
Form	Freeze dried powder
Enzyme Commission Number	EC 1.1.1.47
Activity	250U/mg-solid or more
CAS No.	9028-53-9
Contaminants	NADH oxidase < $1.0 \times 10^{-3}\%$ α -Glucosidase < $1.0 \times 10^{-3}\%$ Glucose-6-phosphate dehydrogenase < $1.0 \times 10^{-3}\%$
Molecular Weight	approx. 101 kDa (Gel filtration)
Isoelectric point	4.5
pH Stability	pH 6.0-7.5 (20°C, 16hr)
Michaelis Constant	NAD*linked: 1.38×10 ⁻² M (D-Glucose) 3.09×10 ⁻⁴ M (NAD*), NADP*linked: 1.25×10 ⁻² M (D-Glucose) 4.07×10 ⁻⁵ M (NADP*)

Fax:1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA



Native Microorganism Glucose Dehyrogenase (NAD(P)-dependent)

Specificity	Specific for ß-D,-Glucose or 2-Deoxy-glucose (Either NAD* or NADP* serves as coenzyme.)
Optimum pH	9
Optimum temperature	55°C
Thermal stability	45°C (15min-treatment with 50mM K-phosphate buffer, pH 7.0)
Stability	Stable at-20°C for at least one year
Inhibitors	Ag+, Hg²+, Monoiodoacetate
Synonyms	Glucose Dehyrogenase; EC 1.1.1.47; beta-D-glucose: NAD(P)+ 1-oxidoreductase; D-glucose dehydrogenase (NAD(P)+)

Fax:1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA