

## Native *Saccharomyces cerevisiae* Alcohol Dehydrogenase

### Product Information

<b>Cat#</b>	NATE-0035
<b>Abbr</b>	ADH, Native ( <i>Saccharomyces cerevisiae</i> )
<b>Alias</b>	ADH
<b>Similar</b>	Alcohol dehydrogenase
<b>Source</b>	<i>Saccharomyces cerevisiae</i>
<b>Description</b>	Alcohol dehydrogenases (ADH) are a group of dehydrogenase enzymes that occur in many organisms and facilitate the interconversion between alcohols and aldehydes or ketones with the reduction of nicotinamide adenine dinucleotide (NAD <sup>+</sup> to NADH). In Humans and many other animals, they serve to break down alcohols that otherwise are toxic, and they also participate in generation of useful aldehyde, ketone, or alcohol groups during biosynthesis of various metabolites. In yeast, plants, and many bacteria, some alcohol dehydrogenases catalyze the opposite reaction as part of fermentation to ensure a constant supply of NAD <sup>+</sup> .
<b>Applications</b>	Alcohol Dehydrogenase from <i>Saccharomyces cerevisiae</i> is used for gel filtration chromatography and as a gel filtration molecular weight marker. It has been used in bioelectrochemical research to investigate the use of diamond nanoparticles as a surface for protein loading.
<b>Form</b>	Solids containing 300 units/mg protein
<b>CAS No.</b>	9031-72-5
<b>Isoelectric point</b>	5.4-5.8
<b>Specificity</b>	The dried enzyme has been stored for several weeks in a vacuum desiccator with little loss in activity. According to experiments described by A. Kornberg, <sup>3</sup> the enzyme can be stored in the frozen state and can be thawed repeatedly without marked loss of activity.
<b>Unit Definition</b>	One unit will convert 1.0 $\mu$ mole of ethanol to acetaldehyde per min at pH 8.8 at 25°C.



**Creative Enzymes**

*Diagnostic Enzymes*

## **Native *Saccharomyces cerevisiae* Alcohol Dehydrogenase**

<b>Optimum pH</b>	8.6-9.0
<b>Storage</b>	-20°C
<b>Inhibitors</b>	Compounds that react with free sulfhydryls, including N-alkylmaleimides and iodoacetamide. Zinc chelator inhibitors, including 1,10-phenanthroline, 8-hydroxyquinoline, 2,2'-dipyridyl, and thiourea. Substrate analogue inhibitors, including $\beta$ -NAD analogs, purine and pyrimidine derivatives, chloroethanol, and fluoroethanol.
<b>Synonyms</b>	aldehyde reductase; ADH; alcohol dehydrogenase (NAD); aliphatic alcohol dehydrogenase; ethanol dehydrogenase; NAD-dependent alcohol dehydrogenase; NAD-specific aromatic alcohol dehydrogenase; NADH-alcohol dehydrogenase; NADH-aldehyde dehydrogenase; primary alcohol dehydrogenase; yeast alcohol dehydrogenase; EC 1.1.1.1